

Evidence for Solar-like Oscillations in Arcturus (α Boo)

Alon Retter, Timothy R. Bedding

School of Physics, University of Sydney 2006, Australia

Derek Buzasi

Department of Physics, 2354 Fairchild Drive, US Air Force Academy, CO 80840, USA

Hans Kjeldsen

Theoretical Astrophysics Center, University of Aarhus, 8000 Aarhus C, Denmark

August 7, 2002

Abstract. Observations of the red giant Arcturus (α Boo) obtained with the star tracker on the Wide Field Infrared Explorer (*WIRE*) satellite during a baseline of 19 successive days in 2000 July-August are analysed. The power spectrum has a significant excess of power at low-frequencies. The highest peak is at $\sim 4.1 \mu\text{Hz}$, which is in agreement with the ground-based radial velocity and photometry study of Belmonte et al. (1990a; 1990b). The variability of Arcturus can be explained by sound waves, but it is not clear whether these are coherent p-mode oscillations.

1. Observations and analysis

After the failure of the main mission of the Wide Field Infrared Explorer (*WIRE*) satellite, launched by NASA in 1999 March, its star tracker was used for asteroseismology of bright objects (Buzasi et al. 2000; Buzasi 2002). Here we report on preliminary analysis of observations of the red giant Arcturus (α Boo), which is the brightest star in the northern hemisphere, and has been observed in velocity several times (Belmonte et al. 1990a; 1990b; Merline 1995).

Nearly 1.2 million 0.1-s exposures were obtained during 19 successive days in 2000 July-August. Fig. 1 presents the light curve. The data from each orbit were binned into a single mean point with a typical error of about 15 ppm. The light curve shows changes on time scales of hours to days.

The upper panel of Fig. 2 displays the amplitude spectrum of the data. There is an excess of power at low frequencies. The highest peak at $4.11 \mu\text{Hz}$ is in agreement with the ground-based radial velocity and photometry study of Belmonte et al. (1990a; 1990b).

Table 1 lists the ten highest peaks in the power spectrum. The typical frequency error is $\sim 0.1 \mu\text{Hz}$. Peak 4 ($0.98 \mu\text{Hz}$) is questionable as the data extend over only ~ 2.5 cycles of it. Fig. 3 presents the






Figure 1. The light curve of α Boo during the 19-d WIRE run in 2002. Each point represents a mean of about 4600 0.1-s exposures obtained during one 96-min orbit. The errors on a single point are smaller than its dimensions in the graph.

amplitude spectrum at the frequency range of 0-20 μHz . The dotted vertical lines show the locations of the ten highest peaks.

2. Discussion

The amplitude and frequency of the power excess in Arcturus are consistent with solar-like oscillations (Kjeldsen & Bedding 1995).

The strongest peaks have a frequency spacing of $\Delta\nu \simeq 0.8 \mu\text{Hz}$, which is in excellent agreement with the expected value of 0.9 ± 0.1 (Kjeldsen & Bedding 1995), but not with the value of $5.0 \mu\text{Hz}$ reported by Belmonte et al. (1990a; 1990b). We note, however, that our value of $\Delta\nu$ is very close to our frequency resolution ($\sim 0.55 \mu\text{Hz}$), which is set by the length of the run. The same was true for the observations by Belmonte et al.

Could granulation noise, rather than oscillations, be responsible for the excess power? This is unlikely, since the variability is also detected in velocity, with an amplitude of $\sim 60 \text{ m/s}$ at $4.3 \mu\text{Hz}$ (Belmonte et al. 1990a; 1990b). The ratio of the photometric amplitude to the velocity




Figure 2. Upper Panel – Amplitude spectrum of the *WIRE* data. There is a significant excess of power at low frequencies. The four highest peaks are at 4.11, 2.26, 3.00 and 0.98 μHz respectively. Second panel – The spectral window (after planting a 4.11- μHz sinusoid in the window function with the same amplitude as in the data). Third panel – The power spectrum after prewhitening the four highest peaks. Lower panel – after prewhitening the eight highest peaks.

Table I. The ten highest frequencies in the amplitude spectrum

Peak	Frequency (μHz)	Semi-amplitude (ppm)	comments
1	4.11	980	
2	2.26	800	
3	3.00	760	
4	0.98	620	questionable
5	5.73	520	
6	8.04	420	
7	4.92	380	
8	10.47	270	
9	7.35	270	
10	16.89	240	




Figure 3. Amplitude spectrum of the *WIRE* data in the range 0-20 μHz . The locations of the ten highest peaks are marked by vertical dotted lines.

amplitude is as expected for sound waves, and about ten times greater than expected for granulation noise.

We conclude that the variations in Arcturus can be explained by sound waves, but it is not clear whether these are coherent p-mode oscillations or perhaps a single mode with a short lifetime. A longer time series is required to distinguish between the two options.

References

- Belmonte J.A., Jones A.R., Palte P.L., Roca Cortes T., 1990a, *Ap&SS*, 169, 77
 Belmonte J.A., Jones A.R., Palte P.L., Roca Cortes T., 1990b, *ApJ*, 358, 595
 Buzasi D., 2002, in Aerts C., Bedding T., Christensen-Dalsgaard J., eds, *ASP Conf. Ser. Vol. 259, Radial and Nonradial Pulsations as Probes of Stellar Physics*, p. 616
 Buzasi D., Catanzarite J., Laher R., Conrow T., Shupe D., Gautier T.N.III, Kreidl T., Everett D., 2000, *ApJ*, 532, L133
 Kjeldsen H., Bedding T., 1995, *A&A*, 293, 87
 Merline W.J., 1995, Ph.D. Thesis, The University of Arizona,